

SINGULAR RIEMANNIAN FOLIATIONS: EXCEPTIONAL LEAVES; TAUTNESS

EVA NOWAK

ABSTRACT. For a singular Riemannian foliation whose leaves are properly embedded, we show in the first part of this article the existence of global tubular neighbourhoods, and we develop a global description of the foliation as stratification by types of leaves.

The second part deals with the further restriction to a foliation without horizontal conjugate points, introduced by Lytchak and Thorbergsson, which in the special case of an isometric group action equals the concept of variationally completeness. Therefrom, we deduce a global geometric description – the focal points of the leaves are exactly the singular points – as well as a topological one: tautness of the leaves.

1. INTRODUCTION

It is a classical result of Bott and Samelson [BS58] that the orbits of a variationally complete group action are taut¹. Isometric group actions are generalized to foliations via the notion of singular Riemannian foliation, see [Mol88]; and Lytchak and Thorbergsson defined in [LTh05] the analog of variational completeness for such foliations via the notion of foliation without horizontally conjugate points. The main aim of this paper is to show that tautness also holds in this more general situation. A basic approach is to generalize the explicit construction of linking cycles done by Bott and Samelson. Pursuing this strategy, we need and find global structural characteristics of the foliations concerned. While singular Riemannian foliations are well understood locally, their global structure can be arbitrarily complicated at least in the non compact case.

After a short introduction, we restrict our interest to properly embedded leaves from Chapter 3 forward. At first, we show the existence of global tubular neighborhoods, see Theorem 3.1, which is the basic tool to reveal global structural results in the following.

Then, we have a look at exceptional leaves, i.e. leaves of highest dimension with non-trivial holonomy. This kind of leaves does not occur too often as the set of specifically regular leaves is open, connected and dense in the manifold (Propositions 3.4 and 3.7). We define the type of a leaf and show that the stratum of exceptional leaves as well as the manifold itself are stratified by the types of leaves.

2000 *Mathematics Subject Classification.* Primary 53C20; Secondary 53C12, 53C40.

Key words and phrases. Singular Riemannian foliations, exceptional leaves, Morse theory, tautness.

¹In the context of foliations there exists a completely different meaning of this notion, see [CC00], but we use *tautness* only in the Morse theoretical meaning as introduced in [GH91] as well as [TTh97].

In the third chapter, we analyse foliations without horizontally conjugate points (with properly embedded leaves). Locally constructed variations of horizontal geodesics can be extended globally, which leads to the geometric characterization formulated in Theorem 4.14: A singular Riemannian foliation has no horizontally conjugate points if and only if the singular points are exactly the focal points of the leaves. For this theorem, we need the extension of homothetical transformations, defined in [Mol88] within tubular neighborhoods, to the negatives which is done in Proposition 4.5. Even more, we show that the homothetical transformations are diffeomorphisms that respect not only the leaves but also the types of leaves. There are two other papers that among other things show the existence of negative homotheties, namely the one of Alexandrino and Töben, [AT07], and the one of Lytchak and Thorbergsson, [LTh07]. As these proofs are contemporaneous but independent and use completely different methods, we include our proof here.

With these structural results we finally can address ourselves to Morse theory of the leaves, i.e. the tautness problem. As a first step, we show that the leaves of a singular Riemannian foliation without horizontally conjugate points are 0-taut. Therefrom, we conclude that there are no exceptional leaves if the foliated manifold is simply connected. At last, we show Theorem 5.4: The leaves of a singular Riemannian foliation without horizontally conjugate points are taut.

Acknowledgements. This paper essentially is my thesis which I wrote under the supervision of G. Thorbergsson at the University of Cologne. He deserves my gratitude for the constantly encouraging support during the last years. Furthermore, I want to thank A. Lytchak and S. Wiesendorf for essential hints towards Proposition 4.9 and Theorem 5.4 in this generality.

2. PRELIMINARIES

2.1. Definitions and notation. Let (M, g) be a Riemannian manifold of dimension n . A decomposition $\mathcal{F}$ of M into smooth, connected, injectively immersed submanifolds, the *leaves*, is called a *singular Riemannian foliation* if the following two conditions hold:

- (I) The modul $\Xi_{\mathcal{F}}$ of smooth vector fields on M that are tangential to the leaves is *transitive*, i.e. for a leaf N_p through p , and for every vector $v \in T_p N_p$ there is a vector field $X \in \Xi_{\mathcal{F}}$ such that $X(p) = v$. Thus, the foliation is *singular* in the sense of [Ste74] and [Su73].
- (II) The metric g on M is *adapted* to $\mathcal{F}$: Each geodesic in M that hits one leaf perpendicularly remains perpendicular to all leaves it passes through. Such a geodesic is called *horizontal*.

Let k denote the maximal dimension of the leaves. A leaf is *regular* if its dimension is equal to k , otherwise it is *singular*. Equally, a point p in M is *regular* (resp. *singular*) if the leaf N_p is regular (resp. singular). The manifold M is stratified by the dimension of the leaves. The *regular stratum* Σ_{reg} is the union of all leaves of dimension k . The union of all leaves of lower dimension, the *singular stratum*, is denoted by Σ_{sing} . Each stratum Σ_q of leaves of dimension q is an embedded submanifold, $\bigcup_{q' \leq q} \Sigma_{q'}$ is closed, and the regular stratum is open, connected, and dense in M .

For the leaf $N_p \subset M$ through p , let νN_p denote the normal bundle of N_p in $TM|_{N_p}$. If $P \subset N_p$ is a relatively compact, open, and connected subset, $B_\varepsilon(P)$ denotes the bundle of open normal balls with radius ε around P in $N_p \subset \nu N_p$.

For ε sufficiently small, the normal exponential map is a diffeomorphism of $B_\varepsilon(P)$ onto its image in M , which is called a *(distinguished) tubular neighborhood of P (resp. around p) with radius ε* :

$$\exp^\perp : \nu N_p \rightarrow M; \exp^\perp(B_\varepsilon(P)) =: \text{Tub}_\varepsilon(P).$$


A *tube* of radius $\delta < \varepsilon$ around P is the union of all points in $\text{Tub}_\varepsilon(P)$ whose distance to P is equal to δ :

$$S_\delta^P := \{x \in \text{Tub}_\varepsilon(P) \mid d(x, P) = \delta\}.$$

Let $\pi : \text{Tub}_\varepsilon(P) \rightarrow P$ denote the projection along horizontal geodesics. We call

$$S_\varepsilon(p) := \pi^{-1}(p)$$

a *slice*. If $y \in S_\varepsilon(p)$, and ε sufficiently small, the leaf N_y cuts the slice transversally.



By the following Lemma 2.1, the connected component of $N_y \cap \text{Tub}_\varepsilon(P)$ that contains y lies in the tube $S_{d(y,P)}^P$. Such a component $P_y \subset N_y$ is called a *plaque*. For more details we refer to [Mol88], Chapter 6, wherefrom we just cite the following facts:

Lemma 2.1. (i) *The projection $\pi : P_y \rightarrow P$ is a surjective submersion, and the fiber $\pi^{-1}(x)$ meets all the plaques in the tubular neighborhood for all $x \in P$.*
(ii) *The distance between neighboring leaves is locally constant for the topology of leaves: $d(\tilde{y}, P) = d(y, P)$ for all $\tilde{y}$ in P_y .*

For every point in Σ_{reg} there exists an open neighborhood U , called *simple*, such that the projection $\pi_U : U \rightarrow \bar{U}$ onto the manifold of plaques is a submersion. The manifold $\bar{U}$ can be endowed with a Riemannian metric $g_{\bar{U}}$ such that the transversal part of g is the pullback of $g_{\bar{U}}$ via π , and

Proposition 2.2. *If γ is a horizontal geodesic in U , its projection $\pi_U(\gamma)$ is a geodesic in $\bar{U}$.*

Let us have a look at a plaque P_N . By the composition of homothetical transformations in the normal bundle νN with $\exp^\perp : B_\varepsilon(P_N) \rightarrow \text{Tub}_\varepsilon(P_N)$, we can define a *homothetical transformation* h_λ with respect to P_N for all positive $\lambda \in \mathbf{R}$ within the domain $\text{Tub}_{\min(\varepsilon, \frac{1}{\lambda}\varepsilon)}(P_N)$, and

Lemma 2.3. *The homothetical transformation h_λ sends plaque to plaque and therefore respects the singular foliation $\mathcal{F}$.*

If γ is a horizontal geodesic, it follows by this lemma that within a tubular neighborhood of $\gamma(0)$

$$\dim(N_{\gamma(t)}) = \text{constant for all } t > 0.$$

2.2. Sliding along the leaves. If $U \subset M$ is simple, the belonging of points to their plaque in U uniquely defines diffeomorphisms between local transversals. Given x and x' in the same plaque, we call the corresponding diffeomorphism *sliding along the leaves from x to x'* , see [Mol88], Section 1.7. By coverings with simple sets, this definition can easily be extended to arbitrary points in a regular leaf and

corresponding local transversals that are sufficiently small. Let c denote a path from x to x' and let h_c denote the germ of this diffeomorphism at x along c . Note that h_c does neither depend on the choice of the covering nor on the choice of local transversals, and concerning c only on its homotopy class. This germ is called *sliding along the leaves from x to x' along c* .

For every point $c(t), t \in [0, 1]$, the slice of minimal radius of the covering tubes $S_{\varepsilon_{\min}}(c(t))$ is a local transversal which then defines for every $y \in S_{\varepsilon_{\min}}(x)$ a unique, continuous path $\tilde{c}_y : [0, 1] \rightarrow L_y$ from y to its image under sliding along c contained in $S_{\varepsilon_{\min}}(x')$, whereby $d(\tilde{c}_y(t), c(t)) = d(y, x)$ for all $t \in [0, 1]$.

Thus, in the following we sometimes speak about sliding of distances instead of points, which can be used not only within regular leaves and simple neighborhoods but – by Lemma 2.1 – always as long as the path c is completely contained in a tubular neighborhood.

2.3. Focal points, cut points, and the cut locus of a leaf. We briefly recall the notions above. For details see [dC93], Section 10.4, and [Tö04]:

Let $N \subset M$ be a submanifold with tangent bundle $TN \subset TM|_N$, normal bundle νN , and normal exponential map $\exp^\perp : \nu N \rightarrow M$. If v is a critical point of $\exp^\perp$, we call v a *focal vector* and $\exp^\perp(v)$ a *focal point*.

Let γ denote a geodesic with $\gamma(0) \in N, \dot{\gamma}(0) \perp N$. An *N -Jacobi field along γ* is a Jacobi field J along γ such that

$$J(0) \in T_{\gamma(0)}N \quad \text{and} \quad J'(0) + A_{\dot{\gamma}(0)}(J(0)) \in \nu_{\gamma(0)}N,$$

where $A_{\dot{\gamma}(0)} : T_{\gamma(0)}N \rightarrow T_{\gamma(0)}N$; $A_{\dot{\gamma}(0)}X = -(\nabla_X^M \dot{\gamma}(0))^T$ denotes the shape operator. J is a N -Jacobi field if and only if it is the variational vector field of a variation f_s of γ , where all paths f_s are geodesics with $f_s(0) \in N, \dot{f}_s(0) \perp N$.

Proposition 2.4. *The following holds for a geodesic γ perpendicular to N in $\gamma(0)$:*

$$\begin{aligned} \gamma(t) \text{ is a focal point of } N &\iff t\dot{\gamma}(0) \text{ is a critical point of } \exp^\perp \\ &\iff \text{there exists a non-vanishing } N\text{-Jacobi} \\ &\quad \text{field } J \text{ along } \gamma \text{ with } J(t) = 0. \end{aligned}$$

The *multiplicity* of $\gamma(t)$ is equal to the dimension of the kernel of $d(\exp^\perp)_{t\dot{\gamma}(0)}$. This also equals the dimension of the space of Jacobi fields along γ vanishing in t .

Let now $N \subset M$ be properly immersed. Let γ_v denote the geodesic starting in the direction of $v \in TM$. Furthermore, let $\nu^1 N$ be the unit normal bundle of N . Then the *cut locus function* is defined by

$$\sigma : \nu^1 N \rightarrow [0, \infty]; \sigma(v) = \sup\{t \in \mathbf{R} \mid d(\gamma_v(t), N) = t\}.$$

The *cut locus of N* is defined to be the set

$$\mathcal{C}_N := \exp^\perp\{\sigma(v)v \mid \sigma(v) < \infty, v \in \nu^1 N\}.$$

We call $v \in \nu N$ *minimal*, if $\|v\| \leq \sigma(\frac{v}{\|v\|})$. The associated geodesic $\gamma_v|_{[0,1]}$ and each of its parametrizations of constant speed are called *minimal geodesics*. A minimal vector v is a *cut vector*, if there is a different minimal $w \in \nu N$ with the same end point; $\exp^\perp(v)$ is then called a *cut point*. Then, $\|v\| = \|w\|$. The following proposition holds:

Proposition 2.5 ([Tö04]). (i) *Each point of $\mathcal{C}_N$ is a cut point or a focal point.*

(ii) *The cut locus function σ is upper semi-continuous, and*

(iii) *if N is properly immersed, σ is continuous and $\mathcal{C}_N$ is closed in M .*

3. EXCEPTIONAL LEAVES – A STRATIFICATION

In general, the leaves of a singular Riemannian foliation are embedded submanifolds. For the following, we additionally assume the embeddings to be proper. It is easy to see that a tubular neighborhood then contains only finitely many plaques of every meeting leaf. As a counterexample take the one-dimensional, irrational foliation of a torus.

3.1. Global tubular neighborhoods. For a singular Riemannian foliation with properly embedded leaves, we can show the existence of global tubular neighborhoods, which will be a useful tool later:

Theorem 3.1. *Let M be complete and $(M, \mathcal{F})$ be a singular Riemannian foliation with properly embedded leaves. Then, for every leaf $N \subset M$ there is an $\varepsilon_N > 0$ such that the normal exponential map*

$$\exp_N^\perp : B_{\varepsilon_N}(N) \rightarrow \text{Tub}_{\varepsilon_N}(N)$$

is a diffeomorphism.


Proof. The proof is divided into two steps:

- (i) The distance of the leaves is globally constant, i.e.: If $\gamma : [0, 1] \rightarrow M$ is a horizontal geodesic that is locally minimizing between N_0 and N_1 , then for every $x \in N_0$ there is a horizontal geodesic of length $L(\gamma)$ from x to N_1 , and this geodesic, too, is locally minimizing between N_0 and N_1 . Hereby, we call γ *locally minimizing*, if there is a neighborhood $P_0 \subset N_0$ around $\gamma(0)$ and a neighborhood $P_1 \subset N_1$ around $\gamma(1)$ such that $d(P_0, P_1) = L(\gamma)$.
- (ii) The distance between a leaf N and its cut locus $\mathcal{C}_N$ is bounded away from zero. For every lower bound $r > 0$, it follows by definition of $\mathcal{C}_N$ that the map

$$\exp_N^\perp : B_r(N) \rightarrow \text{Tub}_r(N)$$

is a diffeomorphism.

We prove step (i) as follows: With regard to x , it is clear that this property is closed. That it is open, too, does not follow directly from the existence of local tubular neighborhoods as, in general, $\gamma(0)$ is not contained in such a neighborhood of a plaque of N_1 , for example if $\gamma(0)$ is singular but N_1 regular. So, we have to show that there is a neighborhood $U \subset P_0$ of $\gamma(0)$ such that $d(x, P_1) = L(\gamma)$ for all $x \in U$. For this purpose take a covering of γ by tubular neighborhoods as follows:



Around the points $\gamma(0)$ and $\gamma(1)$, let the plaques $P_0 \subset N_0$ and $P_1 \subset N_1$ be centers of tubular neighborhoods. The centers of all neighborhoods of the covering shall intersect γ . Furthermore, choose a minimal subdivision

$$0 = t_0 < t_1 < \dots < t_k < t_{k+1} = 1,$$

where P_{t_i} is such a centering plaque for all i , i.e. each point on γ lies either in one or in the intersection of two consecutive tubular neighborhoods. Let $\{s_i\}_{i=1}^{k+1} \subset [0, 1]$ be chosen in such a way that

$$t_0 < s_1 < t_1 < s_2 < \dots < t_k < s_{k+1} < t_{k+1}$$

and $\gamma(s_i)$ lies in the intersection:

$$\gamma(s_i) \in \text{Tub}(P_{t_{i-1}}) \cap \text{Tub}(P_{t_i}).$$

Now, let $U \subset P_0$ be a neighborhood – sufficiently small – of $\gamma(0)$ and $x \in U$. As γ is contained completely in the finite, open covering of tubular neighborhoods, there exists a U so small that the following construction of a broken geodesic is possible within the covering.


In the neighborhood $\text{Tub}(P_0)$, there is a minimal geodesic $c_1 : [0, 1] \rightarrow \text{Tub}(P_0)$, with $c_1(0) = x$ and $c_1(1) \in N_{s_1}$, and the length of c_1 is equal to the length of γ restricted to $[0, s_1]$. Similarly, there is a minimal geodesic c_2 from $c_1(1) \in N_{s_1}$ to P_{t_1} of length $L(\gamma|_{[s_1, t_1]})$. Proceeding like that, we obtain a broken geodesic $c = c_1 c_2 \dots c_{2k+2}$ of length $L(c) = L(\gamma) = d(\gamma(0), P_1)$ which starts in $x \in U$ and ends in P_1 . Thus, the distance between x and P_1 has to be less equal than or equal to $L(c) = L(\gamma)$.

But γ is locally minimizing and equality follows by

$$L(\gamma) \leq d(x, P_1) \leq L(\gamma).$$

To prove step (ii), let $N \in \mathcal{F}$ be arbitrary. Furthermore, choose $\varepsilon > 0$ in such a way that there is a plaque $P_N \subset N$ with tubular neighborhood $\text{Tub}_\varepsilon(P_N)$. In addition, we can assume that the leaf N meets the set $\exp_N^\perp(B_\varepsilon(P_N))$ just once, namely as centering leaf.

Now, let us assume that $\inf\{d(p, \mathcal{C}_N) \mid p \in N\} = 0$. As the leaves are properly embedded submanifolds, Proposition 2.5 tells us that the cut locus function σ_N is continuous. Thus, there exists a leaf L with $d(P_L, P_N) = \frac{\varepsilon}{3}$ where P_L is a connected component of $L \cap \text{Tub}_\varepsilon(P_N)$, there exists a point $q \in N$ with $d(q, \mathcal{C}_N) < \frac{\varepsilon}{3}$, and by (i), there exists a horizontal geodesic γ that realizes the distance $d(q, \mathcal{C}_N)$ and meets the leaf L at length $\frac{\varepsilon}{3}$. But γ cannot be minimal between $q \in N$ and L because the cut locus of N meets γ previous to L . Thus, $\delta := d(q, L) < \frac{\varepsilon}{3}$. Sliding along the leaves with this local distance from L to N up to P_N which is possible by (i) shows that there is a plaque of L lying within the tubular neighborhood $\text{Tub}_\varepsilon(P_N)$ on the δ -sphere of N . But $P_L \subset L$ lies on the $\frac{\varepsilon}{3}$ -sphere around P_N . Once again, sliding along the leaves with distance δ up to P_L shows that there must be one more plaque of N within the neighborhood $\text{Tub}_\varepsilon(P_N)$, a contradiction to the choice of ε !



So, $\inf\{d(p, \mathcal{C}_N) \mid p \in N\} > 0$ has to hold. $\square$

3.2. Holonomy of leaves. In [Mol88], the notion of *leaf holonomy* for Riemannian foliations is defined by sliding along the leaves (see Subsection 2.2 here) on the regular stratum Σ_{reg} :

If c is a loop in x and if we choose the local transversals $T = T'$, then h_c is the germ of a local diffeomorphism with fixed point x . The composition of two loops c_1, c_2 in x is $h_{c_1 c_2} = h_{c_1} \circ h_{c_2}$. Thus, we obtain a group homomorphism

$$h_x : \pi_1(L, x) \rightarrow \text{Diff}_x(T),$$

whose image is the *holonomy group of L in x* . By identification of a local transversal with the quotient $\bar{U}$, the holonomy group is independent of T .

A regular leaf is called *exceptional*, if its holonomy is non-trivial. Furthermore, we call a regular leaf N *specificly regular*, if there is a p in N and a local tubular neighborhood around p wherein every meeting leaf has exactly one plaque. Note that only under the condition of being properly embedded, leaves with trivial holonomy are just the same as specificly regular leaves as the following lemma will show. Thus, we denote this stratum of leaves by Σ_{triv} .

Lemma 3.2. *Let $(M, \mathcal{F}, g)$ be singular Riemannian with properly embedded leaves. Then, a regular leaf N has non-trivial holonomy in x if and only if for every global tubular neighborhood of N there exists a leaf meeting the associated slice $S(x)$ more than once, i.e. if and only if N is not specificly regular.*

To become acquainted with exceptional and specificly regular leaves, we state some easy facts about them previous to the proof of the lemma.

Remark 3.3. (i) For a leaf L , the number of components of the intersection $L \cap S(x), x \in N$, is independent of $x \in N$.

(ii) The holonomy of a leaf is non-trivial in every point of the leaf if it is non-trivial in an arbitrary point.

(iii) For specificly regular leaves, the number of components of the intersection as stated in (i) is constant along connected components of $\Sigma_{\text{triv}} \cap S(x)$.

(iv) The plaques of a leaf B lying in a sufficiently small, global tubular neighborhood of N are lying in *one* tube around N because of globally constant distance.

Proof. The first item results from Theorem 3.1 and locally constant distances of leaves, the second is a direct conclusion of the first. To see the third item, we

consider a tubular neighborhood of a specifically regular leaf B so small that it is contained completely in the global tubular neighborhood of N . Now, combine the local description of a Riemannian foliation with sliding along B to see that the third property is open on $\Sigma_{\text{triv}} \cap S(x)$. Closedness follows since the complement is open. “Sufficiently small” in the fourth item is easily seen to be less than one half of the radius of the global tubular neighborhood (cf. argument (ii) in the proof of theorem 3.1). $\square$

Proof of Lemma 3.2. “ $\Rightarrow$ ” Assume that there is a tubular neighborhood with slice $S(x)$ of x such that every meeting leaf has exactly one plaque in it. As the leaves have the same dimension as N , the intersection of a leaf with the slice is one single point, and sliding along the leaves just produces the identity map.

“ $\Leftarrow$ ” Consider a leaf N with global tubular neighborhood of radius, say, ε ; and let x be a point of N . Further, let B be a leaf with at least two intersection points with the slice $S_\varepsilon(x)$, denote these points by b_1 and b_2 . Sliding along the leaf N of the distance $d(b_1, x)$ to b_2 therefore defines a non-trivial element in the holonomy group of x in N . As B was arbitrary, the claim follows. $\square$

Now, we will show that the exceptional leaves are not as bad as they may seem at a first glance: the stratum Σ_{triv} of leaves of trivial holonomy (i.e. specifically regular leaves) is open, connected, and dense in M ; beyond, the stratum of exceptional leaves $\Sigma_{\text{ex}} = \Sigma_{\text{reg}} \setminus \Sigma_{\text{triv}}$ owns – at least locally – a submanifold structure.

Proposition 3.4. *Let $(M, \mathcal{F})$ be a singular Riemannian foliation with properly embedded leaves. Then, the set of specifically regular leaves is open and dense in M .*

Proof. The stratum of regular leaves Σ_{reg} is open and dense in M . Therefore, it is sufficient to show that Σ_{triv} is open and dense in Σ_{reg} . By Lemma 3.2, the first claim follows.

Let N be exceptional, $x \in N$ and $\varepsilon_0 > 0$ such that $\text{Tub}_{\varepsilon_0}(N)$ is a global tubular neighborhood of N . Let $\varepsilon < \varepsilon_0$ be arbitrary. If every leaf intersected the $\frac{\varepsilon}{3}$ -slice of $x \in N$ only once, N would not be exceptional, so we can find a leaf B intersecting b times. Assume that B is not specifically regular, then iterate the previous consideration as follows:

All the points $\{B_1, \dots, B_b\} = B \cap S_{\frac{\varepsilon}{3}}(x)$ lie in a normal distance sphere of N around x . Around each of these points choose local tubular neighborhoods $\{\text{Tub}_\delta(B_i)\}$ with fixed radius δ in such a way that they do not intersect, they lie within the $\frac{\varepsilon}{3}$ -neighborhood of N , and that each contains just one B_i . Again, have a look at the neighborhoods of one third of this radius. As B is exceptional, there is a leaf C meeting each neighborhood several times and therefore meeting $\text{Tub}_{\frac{\delta}{3}}(B_i) \cap S_{\frac{\varepsilon}{3}}(x)$ more than once, say c times. Thus, there are bc different points of C in the $\frac{\varepsilon}{3}$ -slice of x , and

- (i) the points $C_{i1}, \dots, C_{ic}$ are at the same distance from B_i ,
- (ii) all the bc points $C_{11}, \dots, C_{bc}$ lie on a distance sphere of fixed radius around x .

Let n be the dimension of M and k the dimension of the regular leaves. Then, the dimension of a normal distance sphere of a point in Σ_{reg} is equal to $(n - k) - 1$. The different points of $C \cap S_\varepsilon(x)$ live therefore in an – at most – $(n - k - 2)$ -dimensional set. Thus, the iteration argument has to end after $n - k - 1$ steps which means finding a specifically regular leaf.

The leaf N , the point x , and the radius ε were chosen arbitrarily, so the claim follows. $\square$

To prove that Σ_{triv} is connected, we will use the structure of the exceptional stratum, which can be stated as follows:


Lemma 3.5. *Each connected component of Σ_{ex} is a submanifold of Σ_{reg} with smooth (relative) interior, namely either being an isolated exceptional leaf or locally being the pre-image of totally geodesic, convex sets of the (local) quotient space.*

Remark 3.6. The following proof does not give any information about smoothness of the boundaries of Σ_{ex} in Σ_{reg} resp. in M . It will follow by Lemma 3.11 that every component actually has no boundary (this fact is not used in the meantime).

Proof. Assume that there is a leaf N , to which exceptional leaves accumulate. It is a direct consequence that N itself is exceptional or singular. Let $\text{Tub}_\varepsilon(N)$ be a global tubular neighborhood and let A be exceptional, lying within this neighborhood. Then there exists a horizontal geodesic realising the distance between A and N . By globally constant distance of leaves and by connectedness, we conclude that there is a differentiable strip between the two leaves consisting of minimal geodesics $\{\gamma_a\}_{a \in A}$ of length l . If B is another leaf meeting one of these geodesics at time t , then B meets every such geodesic at time t . This follows by an open-closed argument. To see that B lies totally within this strip, we argue as follows:

Take the distance from $p \in B$ to N as well as the distance from p to A and look at the associated broken geodesic from $\pi_N(p) \in N$ to $\pi_A(p) \in A$. Piecewise, it is of the same length as the geodesics of the strip, so either it must be unbroken (and thus contained in $\{\gamma_a\}_{a \in A}$) or the distance from $\pi_N(p) \in N$ to $\pi_A(p) \in A$ must be less than l , which is impossible.

Lemma 2.3 tells us that B has the same dimension as A . Let $\text{Tub}_\delta(A)$ be a global tubular neighborhood of A and specify B to be a leaf lying within this neighborhood. Choose $a \in A$ arbitrarily and $b \in B$ lying on the unique minimal geodesic from a to N (see the figure below). Within a slice through a , the point b is the only one belonging to B because the distance spheres $S_{d(a,b)}(a)$ and $S_{d(b,N)}(\gamma_a(l))$ intersect only in b . Because A is exceptional, for every neighborhood $U(b) \subset \text{Tub}_\delta(A)$ of b there is a specifically regular leaf which intersects the slice $S_\delta(a)$ more than once.



But locally constant distance forces this leaf to meet $U(b) \cap S_\delta(a)$ several times, otherwise there would be more points of B in $S_\delta(a)$. By definition, B must be exceptional, too.

This consideration holds for every leaf $B \subset \bigcup_{a \in A} \gamma_a \cap \text{Tub}_\delta(A)$. Within Σ_{reg} , the stratum Σ_{ex} is closed. Thus, a boundary leaf $B \in \bigcup_{a \in A} \gamma_a \cap \overline{\text{Tub}_\delta(A)}$ also

is exceptional. Assigning the previous part of A to this boundary leaf B and repeating the proof yields that all leaves of the strip $\bigcup_{a \in A} \gamma_a$ between N and A are exceptional.

In the local quotient space $\bar{U}$, minimal geodesics connecting arbitrary points of $\pi_U(\Sigma_{\text{ex}} \cap U)$ thus lie completely in $\pi_U(\Sigma_{\text{ex}} \cap U)$. This set is locally convex. Therefore, its interior is well-defined and can be endowed with a smooth submanifold structure, see p. 139 of [CE75]. Hence, the preimage $\Sigma_{\text{ex}} \cap U$ of the submersion π_U is smooth in its interior, too. $\square$

Proposition 3.7. *The set Σ_{triv} of leaves with trivial holonomy is connected.*

Proof. Assume $\Sigma_{\text{reg}} \setminus \Sigma_{\text{ex}}$ to be not connected. The component $\mathcal{Z} \subset \Sigma_{\text{ex}}$ that separates the stratum Σ_{triv} has to be $(n-1)$ -dimensional and without boundary. Hence, the fibers of the normal bundle of $\mathcal{Z}$ are one-dimensional.

First case: An exceptional leaf N and thus all regular leaves are of dimension $n-1$. Within every local tubular neighborhood (sufficiently small) of $p \in N$, there is – by lemma 3.2 – a regular leaf having several plaques within a corresponding tube. If N separates the specifically regular stratum, N separates the tube. Transversally, meaning within the quotient, the tube consists of only two points, which then have to belong to different components of Σ_{triv} . But both belong to the same leaf which is connected, a contradiction.

Second case: The component $\mathcal{Z}$ is a union of exceptional leaves. Let $p \in L_p \subset \mathcal{Z}$ and $\text{Tub}_\varepsilon(L_p)$ be a global tubular neighborhood of L_p . By assumption, $\text{Tub}_\varepsilon(L_p) \cap \Sigma_{\text{triv}}$ is separated by $\mathcal{Z}$. Beyond, there is a leaf $N \subset \Sigma_{\text{triv}}$ arbitrarily close to p which intersects $S_\varepsilon(p)$ several times. Choose $x, y \in N \cap S_\varepsilon(p)$, and a path $c : [0, 1] \rightarrow N$ from $x = c(0)$ to $y = c(1)$. At every point $c(t)$, $t \in [0, 1]$, take the distance $d(c(t), \mathcal{Z})$. This is constant along c , hence $d(x, \mathcal{Z}) = d(y, \mathcal{Z})$. As $\mathcal{Z}$ is a $(n-1)$ -dimensional submanifold of Σ_{reg} , the points x and y have to be in different components of $\text{Tub}_\varepsilon(L_p) \cap \Sigma_{\text{triv}}$. But again, both belong to the same leaf, a contradiction. $\square$

By remark (iii) coupled with Propositions 3.4 and 3.7 we conclude:

Corollary 3.8. *In a global tubular neighborhood of an exceptional leaf, all specifically regular leaves intersect a slice equally often (and more than once).* $\square$

3.3. The stratification of Σ_{ex} and M . The last subsection, particularly the last Corollary 3.8, leads to the following, well-defined notion:

Definition 3.9. The *type of an exceptional leaf* N_x is the natural number

$$a_N := \#\{S_\varepsilon(x) \cap L \mid L \subset \Sigma_{\text{triv}} \cap \text{Tub}_\varepsilon(N)\}.$$

Combining the local description of a singular Riemannian foliation with the Tube Theorem 3.1 and Corollary 3.8 yields

Lemma 3.10. *Let N be exceptional of type a_N resp. specifically regular with global tubular neighborhood $\text{Tub}_\varepsilon(N)$. Then, the orthogonal projection*

$$\pi : \text{Tub}_\varepsilon(N) \rightarrow N$$

restricted to a leaf $L \subset \text{Tub}_\varepsilon(N)$ is an $\frac{a_N}{a_L}$ -fold covering resp. a global diffeomorphism. $\square$


A first step towards the stratification shall be the following lemma generalising the claim of Lemma 3.5 as announced. Let $c : (-\varepsilon, \varepsilon) \rightarrow M$ be a horizontal geodesic.

Lemma 3.11. *If $c(0)$ is an accumulation point of both exceptional points and specifically regular points along c , so $c(0)$ must be singular.*

Remark 3.12. This lemma holds for singular Riemannian foliations with properly embedded leaves. Within the restriction to foliations without horizontally conjugate points, we have the stronger Proposition 4.9.

Proof. Without restriction and by Lemma 3.5 we can assume $\varepsilon > 0$ to be so small that $\text{Tub}_\varepsilon(N_0)$ is globally defined and $c|_{(-\varepsilon,0)}$ only consists of exceptional points whereas on $c|_{(0,\varepsilon)}$, there only lie specifically regular points. Furthermore, assume N_0 to be exceptional (by Proposition 3.4, it is clear that N_0 cannot be specifically regular).

Let $x \in \Sigma_{\text{triv}} \cap c$ such that $d(x, c(0)) = \frac{\varepsilon}{4}$. The corresponding leaf N_x meets the tube $\text{Tub}_\varepsilon(N_0)$ more than once, namely a_{N_0} -times (see the following figure).



All intersection points $\{x_i\}$ with the slice through $c(0)$ lie on the distance sphere $S_{\frac{\varepsilon}{4}}^{c(0)}$. We know that $y := c(-\frac{\varepsilon}{4}) \in \Sigma_{\text{ex}}$ and of course $d(x, y) = \frac{\varepsilon}{2}$. Sliding of this distance along the leaf N_x to the different points $\{x_i\}$ yields the fact that the leaf N_y , too, meets the tube $\text{Tub}_\varepsilon(N_0)$ several times. Because of the fixed distance to N_0 , the intersection points $\{y_i\}$ of N_y and the slice of $c(0)$ lie on the sphere $S_{\frac{\varepsilon}{4}}^{c(0)}$. By $d(x, y) = \frac{\varepsilon}{2} = d(x_i, y_i)$ for all i , we conclude, that each y_i is the antipode of the corresponding x_i . But N_y is exceptional, thus we find specifically regular leaves with several plaques within local tubular neighborhoods around the y_i . Choose such neighborhoods so small that neither $c(0)$ nor one of the x_j is contained in it. Hence, a specifically regular leaf that defines the type a_{N_y} must have at least $a_{N_y} a_{N_0}$ plaques within $\text{Tub}_\varepsilon(N_0)$ around N_0 . This contradiction shows that N_0 is singular. $\square$

From the beginning, we have called the set Σ_{ex} a stratum; now we will show that this intuitional naming is correct in the sense of the following common definition.

A locally finite partition of a manifold M into locally closed, connected submanifolds M_i ($i \in I$) of M , the *strata*, is called a *stratification* if the following condition

holds: For every $i \in I$ denote the closure of $M_i \subset M$ by $\overline{M}_i$. Then

$$\overline{M}_i = M_i \cup \bigcup_{j \in I_i} M_j, \text{ wobei } I_i \subset I \setminus \{i\} \text{ and } \dim M_j < \dim M_i \text{ for all } j \in I_i.$$

Remark 3.13. Recall that in the case of a singular Riemannian foliation coming from the orbit decomposition of a proper group action, the subdivision into orbit types is not only a stratification but also fulfills the tougher technical conditions of a Whitney stratification. We conjecture that this is the case in general.

The last argument to do for the stratification by types of leaves as in 3.9 is

Lemma 3.14. *If p is an accumulation point of exceptional points $\{p_i\}_{i=1}^\infty$ and not singular, then N_p is exceptional of type*

$$a_{N_p} \geq \limsup \{a_{N_i}\}, \text{ where } a_{N_i} := a_{N_{p_i}}.$$

Beyond, the function $x \mapsto a_{N_x}$ defined along a horizontal geodesic $c \subset \Sigma_{\text{ex}}$ is constant except for isolated points, i.e. nearly all leaves intersecting c are of the same type, the remaining points are of higher type.

Proof. For every tubular neighborhood of p there is an $i_0 \in \mathbb{N}$ such that the exceptional leaves $\{N_{p_i}\}_{i \geq i_0}$ meet that neighborhood. Additionally, for every $i \geq i_0$ there are specifically regular leaves, meeting the neighborhood a_{N_i} -times, namely those who define the type of N_{p_i} . By assumption, the leaves are properly embedded, therefore the number a_{N_p} and thus the right hand side of the inequality are finite, and the first claim follows.

Now, restrict the attention to a connected subset $U \subset \Sigma_{\text{ex}}$ and define the set of leaves of minimal type, which is open in U . Using the methods of the proof of Proposition 3.4, as there are taking thirds of tubular neighborhoods, sliding of distances, and iteration, we can interpret these leaves as *regular in U* , i.e. they have to meet tubular neighborhoods of exceptional leaves of higher type more than once. Hence, if two different types of leaves accumulate to one point along a horizontal geodesic in Σ_{ex} , we can carry over the Lemma 3.11 directly and conclude that the accumulation point has to be singular, a contradiction. $\square$

With the work already done, the technical Lemma 3.14 can be reformulated as

Proposition 3.15. *The decomposition of Σ_{ex} with respect to the types of the leaves*

$$\Sigma_{\text{ex}} = \dot{\bigcup}_{a_N} \{N \text{ is of type } a_N\}$$

is a stratification. $\square$

Molino has shown that the decomposition of $(M; \mathcal{F})$ with respect to the dimension of the leaves is a stratification. Thus, we conclude with the help of Lemma 3.14 and Proposition 3.15:

Theorem 3.16. *The finer decomposition of a singular Riemannian foliation $(M, \mathcal{F})$ with properly embedded leaves into the sets of specifically regular leaves, strata of types of exceptional leaves, and dimensional strata of singular leaves is a stratification.* $\square$

4. FOLIATIONS WITHOUT HORIZONTALLY CONJUGATE POINTS

The notion of a foliation without horizontally conjugate points was introduced by Lytchak and Thorbergsson in [LTh05] as follows.

Definition 4.1. A singular Riemannian foliation is *without horizontally conjugate points* if the following holds for all leaves N and all geodesics γ perpendicular to N : Every N -Jacobi field along γ that is tangential to a leaf different from N is tangential to every leaf met by γ .

We call a point $q \in M$ *horizontally conjugate to p* if there is a horizontal geodesic $\gamma : [0, 1] \rightarrow M$ with $\gamma(0) = p, \gamma(1) = q$ and an N_p -Jacobi field J along γ such that $J(0) \in T_p N_p, J(1) \in T_q N_q$, but J is not everywhere tangential.

4.1. Tangential Jacobi fields, homothetic transformations. Let $(M, \mathcal{F})$ be a singular Riemannian foliation and let γ be a horizontal geodesic. As in [LTh05] we define the vector space

$$\mathcal{K} := \bigcap_t \{J \mid J \text{ is a } N_t\text{-Jacobi field along } \gamma\},$$

whereby the t 's cover the domain of definition of γ .

We will call a Jacobi field J along γ *everywhere tangential*, if for all t it is tangential to the leaf N_t . Thus, every $J \in \mathcal{K}$ is everywhere tangential.

At least if the foliation has no horizontally conjugate points, we can reverse this implication as follows.

Lemma 4.2. *Let $(M, \mathcal{F})$ be a singular Riemannian foliation without horizontally conjugate points. If J is an N_{t_0} -Jacobi field that is everywhere tangential along γ , then J is an N_t -Jacobi field for all $t \in \mathbf{R}$, i.e. $J \in \mathcal{K}$.*

Remark 4.3. This lemma is a non-mentioned detail of the proof of Proposition 3.1 in [LTh05] which states that, within the notation used above,

$$T_{\gamma(t)} N_t = \{J(t) \mid J \in \mathcal{K}\}.$$

In that article is shown that for every t_0 in the domain of γ and for every $v \in T_{\gamma(t_0)} N_{t_0}$ there is a N_{t_0} -Jacobi field J which is tangential to the corresponding leaf through $\gamma(t)$ for all t . To prove this, a curve $\alpha \subset N_{t_0}$ with $\dot{\alpha}(0) = v$ is lifted to a neighboring leaf B within a tubular neighborhood of $\gamma(t_0)$ via the local Riemannian submersion $\pi_{N_{t_0}}$. The geodesics pointwise connecting both curves define a horizontal variation $f : (-\varepsilon, \varepsilon) \times \mathbf{R} \rightarrow M$ of γ . The appropriate Jacobi field J along γ is tangential twice, namely in N_{t_0} and in B , hence – by assumption – everywhere.

As α lies in N_{t_0} , there actually is a family of Jacobi fields J_s (along the geodesics $f_s(\cdot)$), that are tangential to the leaves $N_{f_s(t)}$ for all t .

Proof of Lemma 4.2. Fix t and define $c : (-\varepsilon, \varepsilon) \rightarrow M$, $c(s) := f(s, t)$, to be a path through the variation f at time t . Then,

$$\dot{c}(0) \in T_{\gamma(t)} N_t \quad \text{and} \quad \dot{c}(s) \in T_{f_s(t)} N_{f_s(t)}.$$

We now verify that – if necessary diminish ε – for all $s \in (-\varepsilon, \varepsilon)$, the geodesics $f_s(\cdot)$ meet the leaf N_t at time t . For this purpose let $s_0 \in (-\varepsilon, \varepsilon)$ be a point with

$$\dim(N_{c(s_0)}) = \max\{\dim(N_{c(s)}) \mid s \in (-\varepsilon, \varepsilon)\}.$$

Without loss of generality consider s_0 to be in the interval $(-\varepsilon, 0)$. There is a neighborhood of s_0 in M containing leaves of the same or higher dimension

only. By maximality of the leaf dimension in $c(s_0)$ along c , there is a $\delta > 0$ with $c|_{(s_0-\delta, s_0+\delta)} \subset \Sigma_{\dim(N_{c(s_0)})}$. Restrict the attention to the stratum $\Sigma_{\dim(N_{c(s_0)})}$ to see that an everywhere tangential path cannot cross several leaves:

For $c(s_0)$, there is a local, centered chart $\psi : U_{s_0} \rightarrow V \times W \subset \mathbf{R}^k \times \mathbf{R}^{n-k}$ such that $\psi(U_{s_0} \cap N_{s_0}) \subset \mathbf{R}^k$, and if N is a leaf with $N \cap U_{s_0} \neq \emptyset$, then

$$\psi(N \cap U_{s_0}) \subset \mathbf{R}^k \times \{\text{isolated points}\}.$$

Denote the composition $\psi \circ c : (s_0 - \delta, s_0 + \delta) \rightarrow V \times W$ by two components (c_1, c_2) . As $\dot{c}(s) \in T_{f_s(s_0)}N_{f_s(s_0)}$, we conclude $\dot{c}_2 \equiv 0$ and $c(s) \in N_{s_0}$ for all $s \in (s_0 - \delta, s_0 + \delta)$.

Let $\tilde{\delta}$ be maximal in the sense that $c|_{(s_0-\delta, s_0+\tilde{\delta})}$ is contained in the leaf $N_{c(s_0)}$. This interval then is open as seen above. We now look at the boundary point $c(s_0 + \tilde{\delta})$: Consider a tubular neighborhood of the leaf $N_{c(s_0+\tilde{\delta})}$, wherein the plaques of the leaves have locally constant distance to $N_{c(s_0+\tilde{\delta})}$. The path $c|_{[s_0, s_0+\tilde{\delta}]}$ is compact, hence c crosses finitely many plaques during this time. Thus, there is a $\rho > 0$ such that the distance from $c(s_0 + \tilde{\delta})$ to $c(s)$ is bigger than or equal to ρ for all $s \in [s_0, s_0 + \tilde{\delta}]$, contradicting continuity of c . Hence, $c|_{(-\varepsilon, \varepsilon)} \subset N_{c(s_0)} = N_{c(0)} = N_t$.

As $f_s(\cdot)$ is perpendicular to N_{t_0} for all s , $f_s(\cdot)$ is perpendicular to N_t . Therefore, f is a N_t -variation, and J is a N_t -Jacobi field. $\square$

The preceding considerations reveal for singular Riemannian foliations

Proposition 4.4. *A curve $c \subset M$ that is always tangential to the leaves, i.e. for all s holds $\dot{c}(s) \in T_{c(s)}N_{c(s)}$, lies in one leaf.*

With regard to Lemma 4.2 and Remark 4.3, we now can define negative homothetical transformations as follows.

Proposition 4.5. *If $(M, \mathcal{F})$ is a singular Riemannian foliation without horizontally conjugate points, then*

- (i) *except at isolated points, $\dim N_{\gamma(t)} = \max\{\dim N_{\gamma(s)} \mid s \in \mathbf{R}\}$ along any horizontal geodesic γ .*
- (ii) *The homothetical transformations h_λ also exist for negative λ and respect the foliation. Particularly, the reflection at a leaf respects the foliation.*

Proof. To prove the first item, let $\gamma(t_0)$ be such that the leaf through $\gamma(t_0)$ is of maximal dimension (along the geodesic); let $N_{\gamma(t_1)}$ be of smaller dimension. By maximality, we can choose t_1 in such a way that $\gamma|_{[t_0, t_1]} \subset \Sigma_{\dim(N_{t_0})}$. Now consider a local tubular neighborhood of $\gamma(t_1) \in N_{t_1}$ of normal radius ρ and its intersection with the stratum $\Sigma_{\dim(N_{t_1})}$. Following [Mol88], p. 196, this is a union of geodesic arcs of length 2ρ perpendicular to N_{t_1} . As γ is perpendicular to N_{t_1} in $\gamma(t_1)$, too, we have

$$\gamma|_{(t_1-\rho, t_1+\rho)} \subset \Sigma_{\dim(N_{t_1})} \text{ or } (\gamma|_{(t_1-\rho, t_1+\rho)} \setminus \{\gamma(t_1)\}) \cap \Sigma_{\dim(N_{t_1})} = \emptyset,$$

i.e. the geodesic arc $\gamma|_{(t_1-\rho, t_1+\rho)}$ is either contained completely in the stratum $\Sigma_{\dim(N_{t_1})}$ or not at all except the point $\gamma(t_1)$.

The first inclusion cannot be fulfilled as $\gamma|_{[t_0, t_1]} \subset \Sigma_{\dim N_{t_0}}$ and we assumed the strata to be different. Because of the existence of positive homothetical transformations, it is only left to show that there is an $\varepsilon < \rho$, such that $\gamma(t_1 + \varepsilon) \in \Sigma_{\dim(N_{t_0})}$.

Denote the maximal dimension of leaves along γ by k , and the dimension of N_{t_1} by l . The intersection of a slice through $\gamma(t_1)$ with a plaque of a leaf $N_{t_1-\varepsilon}$

is $(k - l)$ -dimensional and lies within a $(\dim(M) - l)$ -sphere around $\gamma(t_1)$. All geodesics beginning in $\gamma(t_1)$ and ending in $N_{t_1-\varepsilon}$ at the same time as γ form a $(k - l)$ -dimensional variation of horizontal geodesics which can be continued up to infinity. The negative starting vectors are of course tangential to the negatives of the curves and span a $(k - l)$ -dimensional submanifold in $T_{\gamma(t_1)}M$. Within the tubular neighborhood, we can assume the existence of unique geodesics between arbitrary points, and the negative variation up to $t_1 + \varepsilon$ has to stay $(k - l)$ -dimensional. Hence, there is only one leaf meeting the variation at time $t_1 + \varepsilon$ and this is k -dimensional which means maximal.

The second item now is easily seen: The proof of (i) shows the existence of the homothetical transformation h_{-1} at N_{t_1} . Compose this with positive ones. $\square$

Corollary 4.6. *Let N be a leaf in M with global tubular neighborhood $\text{Tub}_\varepsilon(N)$. Then, the homothetical transformations*

$$h_\lambda : S_\delta^N \rightarrow S_{\lambda\delta}^N$$

(as long as $\lambda \in (-\varepsilon, \varepsilon)$ and δ are defined) map leaves diffeomorphically onto each other.

Proof. Consider a leaf L_1 . By Proposition 4.5, h_λ locally maps open sets of L_1 diffeomorphically to open sets of other leaves. The image $h_\lambda(c)$ of an arbitrary path $c : [0, 1] \rightarrow L_1$ is tangential to the leaves as a path through the variation of horizontal geodesics defining h_λ . Hence, h_λ maps L_1 into one leaf L_2 . Further, $h_\lambda : L_1 \rightarrow L_2$ is injective.

Assume that there is $x \in L_2$ contained in the closure of $h_\lambda(L_1)$ but not in its image. Then there is a variation through horizontal geodesics whose endpoints in L_2 converge to x . The limit curve is a horizontal geodesic γ beginning in L_1 and meeting N , too. Thus, $\gamma(0) \in L_1$ is a preimage of x with respect to h_λ which means that $h_\lambda(L_1)$ is open and closed in L_2 , hence equal to L_2 . $\square$

Remark 4.7. In [AT07] and [LTh07] is shown (by completely different methods) that the reflection respects the foliation in general, i.e. without the assumption of no horizontally conjugate points. By Theorem 3.1 we thus generalize: *For a singular Riemannian foliation with properly embedded leaves, the homothetical transformations defined within global tubular neighborhoods are diffeomorphisms between the leaves.*

Remark 4.8. Due to an essential hint of A. Lytchak, we can specify the last corollary even to the types of leaves, which is stated in the next proposition. Thus, if the foliation has no horizontally conjugate points, a situation as in Lemma 3.11 cannot occur which means: If there lie specifically regular points on a horizontal geodesic, singular points as well as exceptional points are isolated along this geodesic, and Lemma 3.14 does not only hold in Σ_{ex} but in M itself.

Proposition 4.9. *If $(M, \mathcal{F})$ is a singular Riemannian foliation without horizontally conjugate points and with properly embedded leaves, then the homothetical transformations respect the types of leaves, i.e. they map a leaf N of type a_N diffeomorphically onto a leaf of the same type.*

Proof. As the leaves are properly embedded, the foliation is closed. We can then use Theorem 1.7 of [LTh07] to see that the quotient $M/\mathcal{F}$ is isometric to a good

Riemannian orbifold N/Γ without conjugate points. The manifold N and the discrete group Γ are constructed in such a way that the stratification by types of leaves of M determines the group Γ of isometries of N (for details see [LTh07]). Because of our Lemma 3.14, it only remains to check whether the reflection, i.e. the homothetical transformation h_{-1} , at a singular leaf respects the types of the neighboring leaves. And this is true on M because it is an obvious fact on N stratified by the action of Γ . $\square$

We close this subsection by citing a lemma proven by Lytchak and Thorbergsson in [LTh07], Subsection 4.5, which we will need later.

Lemma 4.10. *Let $(M, \mathcal{F})$ be a singular Riemannian foliation, $\gamma : (-\infty, \infty) \rightarrow M$ a horizontal geodesic. Denote by W^γ the space of horizontal Jacobi fields along γ , i.e. all Jacobi fields that come from variations of γ through horizontal geodesics in which all geodesics meet the same leaves at the same time. For every $t \in (-\infty, \infty)$, W^γ then spans the tangential space of the leaf N_t :*

$$\{J(t) \mid J \in W^\gamma\} = T_{\gamma(t)}N_t.$$

In particular, $J(\tilde{t}) \in T_{\gamma(\tilde{t})}N_{\tilde{t}}$ for all $\tilde{t}$.

4.2. Singular points and focal points. The preceding section already points at the possibility to characterise singular points locally by variations through horizontal geodesics which leads to the notion of focal points. In this chapter, we will now have a deeper view onto the relation of these attributes.

For the following, let $(M, \mathcal{F})$ be a singular Riemannian foliation without horizontally conjugate points; let γ be a horizontal geodesic in M meeting the leaf N_0 at time 0. Furthermore, consider N_0 to be of maximal dimension along γ . Then,

Lemma 4.11. $\gamma(t)$ is a focal point of $N_0 \iff \dim N_t < \dim N_0$.

Proof. $\Leftarrow$: This can be seen in the proof of proposition 4.5: For a leaf $N_{t+\varepsilon}$ near $\gamma(t)$, we find a $(k-l)$ -dimensional variation of horizontal geodesics starting in $\gamma(t)$ that is everywhere tangential. A singular point on γ thus is a focal point to every leaf of higher dimension met by γ .

$\Rightarrow$: Let $f(s, \tilde{t})$ denote a non-trivial variation of horizontal geodesics starting in N_0 and meeting the leaf N_t in $\gamma(t)$. The corresponding Jacobi field is tangential in $\gamma(0)$ and in $\gamma(t)$, hence everywhere. Therefore, $f(s, \tilde{t})$ is an $N_{\tilde{t}}$ -variation for every leaf met by γ at time $\tilde{t}$ with focal point $\gamma(t)$, if $N_{\tilde{t}}$ is of higher dimension than N_t . Again, take a tubular neighborhood of $\gamma(t) \in N_t$ so small that there exist unique geodesics between arbitrary points, and let ε be such that $\gamma(t + \varepsilon)$ is contained in it. Then, $f(s, t + \varepsilon) \in S_\varepsilon^{\gamma(t)}$ for all s , where $S_\varepsilon^{\gamma(t)}$ denotes the normal sphere of radius ε around $\gamma(t)$. Assuming N_t to have maximal dimension now leads to a contradiction: The leaf $N_{t+\varepsilon}$ must have the same dimension and

$$N_{t+\varepsilon} \cap S_\varepsilon^{\gamma(t)} = \text{finitely many points}$$

while the variation is non-trivial and its geodesics are unique, so $\dim N_t < \dim N_0$. $\square$

Remark 4.12. In a singular Riemannian foliation (without additional conditions), singular points are easily seen to be focal points of leaves nearby. But in general, there are focal points in regular leaves, too! A trivial example is the sphere S^2 with the standard metric, foliated by points. Geodesics are always horizontal, and

conjugate points are horizontally conjugate. Obviously, they are focal points to each other, and regular by definition.

With regard to the last proof and the proof of Proposition 4.5, we can improve Lemma 4.11 as follows.

Proposition 4.13. *For every singular point p in M and every regular leaf N there is a geodesic $\gamma : [0, 1] \rightarrow M$ with $\gamma(0) =: q \in N, \gamma(1) = p$, that is perpendicular to $N = N_q$, and p is a focal point of N_q along γ of multiplicity*

$$\nu(p) = \dim(N_q) - \dim(N_p).$$

In particular, the multiplicity of p is independent of the choice of γ .

Proof. We just drop a perpendicular from p to N ; let $\gamma : [0, 1] \rightarrow M$ be a geodesic beginning in $q = \gamma(0)$ perpendicularly to $N = N_q$, ending in p . As in the proof of Lemma 4.11, there is a local variation of γ of maximal dimension $\nu(p)$ which can be extended up to the leaf N_q . If this variation was not $\nu(p)$ -dimensional in N_q the point q would be a focal point along γ (namely a focal point of the leaf $N_{t+\varepsilon}$ in the notation used above) which contradicts the regularity of N_q . $\square$

Combining the latest findings with the general form of Proposition 4.5 part (i) as proven in [LTh07] and [AT07], yields the main theorem of this chapter, the geometric characterisation of foliations without horizontally conjugate points.

Theorem 4.14. *A singular Riemannian foliation has no horizontally conjugate points if and only if the singular points are exactly the focal points of leaves.*

To prove the if-part, we will need the following technical lemma that holds under the assumption of singular points being the focal points.

Lemma 4.15. *Let $\gamma : (-\infty, \infty) \rightarrow M$ be a horizontal geodesic. If we have $a < b$ such that $\gamma(b)$ is horizontally conjugate to $\gamma(a)$ along γ , for every $\varepsilon > 0$ (sufficiently small) there is a $\delta > 0$ and $c \in (a + \delta, b + \varepsilon)$ such that $\gamma(c)$ is horizontally conjugate to $\gamma(a + \delta)$.*

Proof. We define $\mathcal{T} := \{J \mid J \text{ is a Jacobi field along } \gamma, J(\tilde{t}) \in T_{\gamma(\tilde{t})}N_{\tilde{t}} \forall \tilde{t}\}$. In analogy to [Wi06] and [LTh05], we can define a bundle $\mathcal{V}_\gamma$ over $\mathbf{R}$ by the following extension of $\mathcal{T}$ at singular points:

$$\mathcal{V}_t = \{J(t) \mid J \in \mathcal{T}\} \oplus \{J'(t) \mid J \in \mathcal{T}, J(t) = 0\}.$$

Then, $\{J'(t) \mid J \in \mathcal{T}, J(t) = 0\} = 0$ if t is regular by the assumption that singular points are focal points. As we know the latter to be isolated along γ , we see that $\mathcal{V}_\gamma$ is a smooth bundle, and $\dim(\mathcal{V}_t) = \dim(T_{\gamma(t_0)}N_{t_0})$ for all t whereby $\gamma(t_0)$ is regular (see Proposition 3.2 in [LTh05]). Define the horizontal part $\mathcal{H}_t$ to be the orthogonal complement of $\mathcal{V}_t$ in $T_{\gamma(t)}N_t$ and denote the bundle by $\mathcal{H}_\gamma$.

Wilkings showed that there is an associated Morse Sturm differential equation to $\mathcal{H}_\gamma$ in $\mathbf{R}^k$, such that its space of solutions $\mathcal{L}$ determines just the horizontal Jacobi fields along γ which vanish in a . By the generalized index theorem of Morse, the corresponding index form

$$I_{a,b}(V, W) = \int_a^b \langle V', W' \rangle - \langle \mathcal{R}(t)V, W \rangle dt$$

has

$$\text{Ker}(I_{a,b}) = \{J|_{[a,b]} \mid J \in \mathcal{L}, J(b) = 0\} \quad \text{and} \quad \text{Ind}(I_{a,b}) = \sum_{t \in (a,b)} \nu(t).$$

Here, V, W lie in $H_0^1([a, b], \mathbf{R}^k)$, the space of absolutely continuous, $\mathbf{R}^n$ -valued maps of $[a, b]$ with square integrable derivative and boundary condition $V(a) = V(b) = 0$; $\mathcal{R}$ is continuous and $\mathcal{R}(t)$ is self-adjoint for all t (here, we use the notation of Lytchak and Thorbergsson in [LTh05] which we refer to for details).

Without loss of generality let $\gamma(b)$ be the first conjugate point to $\gamma(a)$. For every $\varepsilon > 0$ sufficiently small, let $\mathcal{U} \subset H_0^1([a, b + \varepsilon], \mathbf{R}^k)$ be a maximal subspace on which the index form is negative definite

$$I_{a,b+\varepsilon}(V, V) < 0 \quad \text{for all } V \in \mathcal{U}.$$

The Morse Sturm differential equation and thus the index form depend continuously on the end points of the interval. Beyond, there is a neighborhood around $N_{\gamma(a)}$ that contains leaves of same or higher dimension only, so by the index theorem the following holds for all $\delta > 0$ sufficiently small:

$$I_{a+\delta,b+\varepsilon}(\tilde{V}, \tilde{V}) < 0 \quad \text{for all } \tilde{V} \in \tilde{\mathcal{U}},$$

where $\tilde{\mathcal{U}} \subset H_0^1([a + \delta, b + \varepsilon], \mathbf{R}^k)$ with boundary condition $V(a + \delta) = V(b + \varepsilon) = 0$ is a subspace of dimension $\dim(U)$. Thus, continuity implies

$$\exists_{c \in (a+\delta, b+\varepsilon)} : \gamma(c) \text{ is horizontally conjugate to } \gamma(a+\delta). \quad (*)$$

□

Remark 4.16. We can state $(*)$ more precisely: For $\gamma(a + \delta)$ there is a horizontally conjugate point $\gamma(c)$ with $c \in [b, b + \varepsilon]$.

By assumption, there is no horizontally conjugate point between $\gamma(a)$ and $\gamma(b)$. For all $\varepsilon > 0$ and all $W \in H_0^1([a, b - \varepsilon], \mathbf{R}^k)$, we therefore have $I_{a,b-\varepsilon}(W, W) > 0$. If δ in $(*)$ is chosen to be sufficiently small, continuity implies $I_{a+\delta,b-\varepsilon}(W, W) > 0$. Hence, $\gamma(a + \delta)$ has no horizontally conjugate point on $\gamma|_{[a+\delta, b]}$.

Proof of Theorem 4.14. With regard to Lemma 4.11 it remains to show that if focal points are singular, then the foliation is without horizontally conjugate points. We prove this by contradiction.

Assume that there are horizontally conjugate points, i.e. there is a horizontal geodesic $\gamma : [0, 1] \rightarrow M$, points $t_0, t_1 \in [0, 1]$ and a N_0 -Jacobi field J along γ such that $J(t_0) \in T_{\gamma(t_0)}N_{t_0}$, but $J(t_1) \notin T_{\gamma(t_1)}N_{t_1}$. By Lemma 4.10, there exists a N_0 -Jacobifield $\tilde{J}$ along γ that is everywhere tangential and $\tilde{J}(t_0) = J(t_0)$. The difference $J - \tilde{J}$ is a N_0 -Jacobi field that vanishes in t_0 but not identically. Hence, $\gamma(t_0)$ is a focal point of N_0 .

Wilkings shows in [Wi06], Section 3, that then $(J - \tilde{J})'(t_0) \in \nu_{\gamma(t_0)}N_{t_0}$, i.e. $J - \tilde{J}$ is a N_{t_0} -Jacobi field. Now, use lemma 4.10 again: There is $\hat{J}$ such that $J - \tilde{J} - \hat{J}$ is a non-trivial N_{t_0} -Jacobi field vanishing in 0. Hence, $\gamma(0)$ is a focal point of N_0 and thus singular.

By Lemma 4.15, we can disturb horizontally conjugate points, so they are not isolated. But for every $\delta > 0$ (notation as in 4.15), $\gamma(0)$ as well as $\gamma(0 + \delta)$ lie in singular leaves and thus have to be isolated along γ , a contradiction! □

Corollary 4.17. *A singular Riemannian foliation with regular leaves of codimension one has no horizontally conjugate points.*

Proof. Let $N \in \mathcal{F}$ be arbitrary. Bolton shows in [?] that the critical points of the normal exponential map $\exp_N^\perp$ are just the singular points if the foliation is of codimension one. $\square$

5. TAUTNESS OF THE LEAVES

5.1. Taut immersions. In order to draw topological conclusions for the leaves from their geometry we will use Morse theory. At first, let us recall some notation.

A proper embedding $N \rightarrow M$ is called *taut* (with respect to $\mathbf{Z}_2$), if the energy functional $E_p : P(M, N \times p) \rightarrow \mathbf{R}$, $c \mapsto \int_0^1 \|\dot{c}(t)\|^2 dt$, is $\mathbf{Z}_2$ -perfect for every p in M , that is not a focal point of N , where

$$P(M, N \times p) = \{c : [0, 1] \rightarrow M \text{ in } H^1 \mid (c(0), c(1)) \in (N, p)\}$$

denotes the Hilbert space of H^1 -paths starting in N and ending at p , see [GH91] and [TTh97]. It is well known that the energy functional is a Morse function whose critical points in $P(M, N \times p)$ are exactly the geodesics starting perpendicularly to N and parametrized proportionally to arc length. Now, Morse's index theorem tells us that the index of such a geodesic γ is equal to the sum of multiplicities of the focal points along γ .

If $\mathcal{N}_p^r = \{c \in P(M, N \times p) \mid E_p(c) \leq r\}$ denotes the sublevel set of paths with energy less than or equal to $r \in \mathbf{R}$, then the number μ_k^r of critical points of index k of E_p in $\mathcal{N}_p^r$ is finite. Let b_k^r be the k -th Betti number of $\mathcal{N}_p^r$ with respect to $\mathbf{Z}_2$ coefficients, so for all r we can state the weak Morse inequalities

$$\mu_k^r \geq b_k^r \text{ for all } k.$$

A Morse function is called $\mathbf{Z}_2$ -perfect, if equality holds for all r and all k :

$$\mu_k^r = b_k^r.$$

If N is taut, the focal points of N along a horizontal geodesic determine the $\mathbf{Z}_2$ -homology of the space $P(M, N \times p)$ completely.

To show tautness of an immersion, it is a well working strategy to construct linking cycles explicitly, as done by Morse [Mor34], p. 244f, and in the famous paper of Bott and Samelson [BS58] (for a good exposition of this theory see [PT89], for the construction of such cycles in special cases see also [STh71], [GTh02], and [Th83]). As we will do so, too, let us shortly review this construction: The Morse inequalities reveal the deep connection between the k -th homology group of $\mathcal{N}_p^r$ and the geodesics of energy r with index k . Given such a geodesic, we need to construct an appropriate, non-trivial class, the linking cycle, in $H_k(\mathcal{N}_p^r, \mathbf{Z}_2)$ out of the data of the geodesic, i.e. focal points and their multiplicities, to obtain equality.

As above we define

$$\mathcal{N}_p^{r-} = \{c \in P(M, N \times p) \mid E_p(c) < r\}.$$

It is a well known fact that $\mathcal{N}_p^r$ is homotopy equivalent to $\mathcal{N}_p^{r-}$ if r is not critical. And if r is critical, there is an $\varepsilon > 0$, such that

$$H_k(\mathcal{N}_p^r, \mathcal{N}_p^{r-\varepsilon}) = \mathbf{Z}_2^i,$$

where i denotes the number of critical points of E_p with index k and value r . Then the long exact homology sequence tells us that E_p is perfect iff

$$H_k(\mathcal{N}_p^r) \rightarrow H_k(\mathcal{N}_p^r, \mathcal{N}_p^{r-\varepsilon}) \quad (*)$$

is surjective for all k .

Let γ be a critical point of E_p with value r , let

$$1 > t_1 > \dots > t_k > t_{k+1} = 0$$

such that $\{\gamma(t_i)\}_{i=1}^k$ are just the focal points of N along γ . Then

$$\text{ind}(\gamma) = \sum_{i=1}^k \nu(\gamma(t_i)).$$

To show surjectivity in $(*)$ it suffices to construct for every γ a compact, $\text{ind}(\gamma)$ -dimensional manifold K , and a smooth map

$$\lambda : K \rightarrow P(M; N \times p),$$

whereby the following conditions hold:

- (1) $\lambda(x_0) = \gamma$ for exactly one $x_0 \in K$, and there is a neighborhood U of γ such that $\lambda|_{\lambda^{-1}(U)}$ is an immersion,
- (2) γ is the only critical point of E_p in the image of λ , and
- (3) $E_p(\lambda(x)) \leq E_p(\gamma) = r$ for all $x \in K$.

By the negative gradient flow of E_p , the image $\lambda(K)$ can then be deformed into a smooth cycle Z_γ in $P(M, N \times p)$, which represents a non-trivial class in $H_{\text{ind}(\gamma)}(\mathcal{N}_p^r)$. Conditions (i) and (ii) guarantee that the image of this class in $(*)$ is non-trivial, too. So, for every γ we would have found a linking cycle Z_γ , which shows that $H_k(\mathcal{N}_p^r) \rightarrow H_k(\mathcal{N}_p^r, \mathcal{N}_p^{r-\varepsilon})$ is surjective for every k . Thus, tautness of N follows.

5.2. Topology of singular Riemannian foliations without horizontal conjugate points. Let N denote a leaf and $P(M, N \times p)$ the corresponding space of paths ending in a point p that is not a focal point of N . Critical points of the energy integral E_p are horizontal geodesics from N to p with index equal to the sum of focal points counted with multiplicities. Hence, the easiest case for which we have to construct a linking cycle is a geodesic with only one focal point that is of multiplicity one. If all such critical points of index one are completable, we call N *0-taut*. As a first step towards the general construction we show

Theorem 5.1. *Let $(M, \mathcal{F})$ be a singular Riemannian foliation without horizontal conjugate points on a complete Riemannian manifold M . Then, the leaves are 0-taut.*

Remark 5.2. The following construction of a linking cycle obviously also works in the situation of a horizontal geodesic with only one focal point whose multiplicity is arbitrary.

Proof. At first, let N be a regular leaf and let $\gamma : [0, 1] \rightarrow M$ be a critical point of E_p as above, i.e. a horizontal geodesic with exactly one focal point $\gamma(t_0)$ of multiplicity one. By Theorem 4.14, $\gamma(t_0)$ is singular. Hence, by Lemma 3.11 and Lemma 3.14, we can conclude that except isolated points, all points along $\gamma|_{(0, t_0]}$ are of the same type. Furthermore, the isolated points are of higher type.

Choose a global tubular neighborhood $\text{Tub}_\varepsilon(N_{t_0})$ and $\delta < \varepsilon$ such, that $\gamma(t_0 - \delta)$ is specifically regular resp. of minimal type along γ . Then, $N_{\gamma(t_0 - \delta)} \cap S_\varepsilon(\gamma(t_0))$ is a

compact, properly embedded submanifold of dimension $\nu(\gamma(t_0)) = 1$ of $N_{\gamma(t_0-\delta)}$. If it is not connected, we choose the component – topologically a circle – that contains $\gamma(t_0 - \delta)$ and call it K . Now, define

$$f : K \times [0, t_0] \rightarrow M, \text{ with } \begin{cases} f(s, t_0) = \gamma(t_0) \text{ for all } s \in K \\ \text{and } K = \{f(s, t_0 - \delta) \mid s \in K\}, \end{cases}$$

to be the one-dimensional, smooth variation of horizontal geodesics of the same length as $\gamma|_{[0, t_0]}$. This variation is unique because the paths $f_s|_{[t_0-\delta, t_0]}$ are minimal from K to $\gamma(t_0)$ in M for all s .

The path through the variation at time $t_0 - \delta$ is tangential to the leaf $N_{\gamma(t_0-\delta)}$. Obviously, $\frac{\partial}{\partial s} f(s, t_0)$ is tangential to N_{t_0} for all s . Thus,

$$\frac{\partial}{\partial s} f(s, t) \in T_{f_s(t)} N_{\gamma(t)} \text{ for all } s \text{ and all } t;$$

and by proposition 4.4 we have $f(s, 0) \in N$ for all s . Moreover, the set

$$\{f(s, 0) \mid s \in K\} \subset N$$

is a compact submanifold (possibly not properly embedded) of N as image of K :

Cover $\gamma|_{[0, t_0-\delta]}$ by global tubular neighborhoods similar to the covering in the proof of Theorem 3.1. For this purpose, let the centers be the leaf N as well as all the leaves of higher type than $N_{\gamma(t_0-\delta)}$ and, if these tubular neighborhoods do not yet cover $\gamma|_{[0, t_0-\delta]}$, some specifically regular leaves (resp. leaves of the same type as $N_{\gamma(t_0-\delta)}$). Thus, each point $f(s, t)|_{[0, t_0-\delta]}$ lies within exactly one neighborhood or within the section of two consecutive neighborhoods. Denote the centering leaves by $N = N_0, N_1, \dots, N_r \neq N_{\gamma(t_0-\delta)}$ with $N_{\gamma(t_0-\delta)} \subset \text{Tub}(N_r)$. The homotheties defined within the global neighborhoods are diffeomorphisms by Corollary 4.6, and with the help of these maps, we move K along the geodesics $f_s(t)$ from $N_{\gamma(t_0-\delta)}$ to a leaf (of the same type) in the intersection $\text{Tub}(N_r) \cap \text{Tub}(N_{r-1})$, then to a leaf in $\text{Tub}(N_{r-1}) \cap \text{Tub}(N_{r-2})$, etc. up to a leaf L in the intersection $\text{Tub}(N_1) \cap \text{Tub}(N_0)$. The image of K in L , i.e. the set $\{f(s, \tilde{t}) \mid s \in K, \gamma(\tilde{t}) \in L\}$, is mapped to $\{f(s, 0) \mid s \in K\}$ via the orthogonal projection $\pi_N|_L : L \rightarrow N$ in the last step, hence we loose embeddedness in general.

Now define an injective, smooth map $\lambda : K \rightarrow P(M, N \times p)$ by

$$\lambda(s)(t) = \begin{cases} f_s(t), & t \in [0, t_0], \\ \gamma(t), & t \in [t_0, 1]. \end{cases}$$

The compact image $\lambda(K)$ is an injective, smooth cycle of dimension $\nu(\gamma(t_0))$ in $\mathcal{N}_p^{E(\gamma)}$ consisting of broken geodesics of same length and thus of same energy. The only non-broken geodesic is γ . Hence, the critical point γ is completable.

For singular leaves, we can use an idea based on Proposition 2.7 in [TTh97], which in the general case of foliations is due to S. Wiesendorf, see [Wi09]: By definition, a leaf N is taut, if the energy functional E_p is perfect for p not focal, but by a limiting argument, it suffices to show perfectness for all p that are regular. Thus, along a horizontal geodesic, nearly all points are regular. To show that a singular leaf N is taut, we construct a linking cycle $\lambda(K)$ of a regular leaf L nearby and then add to every broken geodesic in $\lambda(K)$ a minimal geodesic connecting the endpoint of the geodesic in $\lambda(K)$ with N . The corresponding map

$$P(M; L \times p) \rightarrow P(M; N \times p); c \mapsto \tilde{c} = \gamma_{c(0)} c$$

is injective and preserves the indices of geodesics as critical points of E_p . Furthermore, the energy levels will just be shifted by a constant. The new set of broken geodesics, $\widetilde{\lambda(K)}$, by construction can be deformed into a linking cycle of N . $\square$

Töben has shown in [Tö04] that a singular Riemannian foliation with properly embedded leaves and with flat sections in a simply connected symmetric space does never have exceptional leaves. As an application of the previous theorem, we generalize this to

Proposition 5.3. *If M is a complete, simply connected manifold and if $(M, \mathcal{F})$ is singular Riemannian without horizontal conjugate points and with properly embedded leaves, there are no exceptional leaves in $\mathcal{F}$.*

Proof. Let N be a leaf and $p \in M$. There is the fibration

$$P(M, N \times p) \rightarrow N; c \mapsto c(0),$$

where $c : [0, 1] \rightarrow M$ is a path with $c(0) \in N, c(1) = p$. The fiber $\Omega_{x_0 p}(M)$ over $x_0 \in N$, that is the space of paths from x_0 to p , is homotopy equivalent to $\Omega_{x_0}(M)$ which is connected as M is simply connected. The long exact sequence of homotopy groups of fibrations yields that $P(M, N \times p)$ is connected:

$$\begin{array}{ccccccc} \cdots & \longrightarrow & \pi_1(N) & \longrightarrow & \pi_0(\Omega_{x_0}(M)) & \longrightarrow & \pi_0(P(M, N \times p)) \longrightarrow \pi_0(N) \longrightarrow 0 \\ & & & & \parallel & & \parallel \\ & & & & 0 & & 0 \end{array}$$

Assume that there is an exceptional leaf $A \subset M$. Then let B be a leaf with trivial holonomy meeting a tubular neighborhood of $x \in A$ several times. Under the boundary condition $B \times x$, the energy integral thus must have at least two local minima. Now use the minimax principle:

There is a critical point of index one, i.e. a geodesic c from B to x with just one focal point, and this critical point connects two components of $\mathcal{B}_x^{E(c)-}$ in $\mathcal{B}_x^{E(c)}$. By Theorem 5.1, there is a one-dimensional linking cycle Z , topologically a circle, in $P(M, B \times x)$ with

$$E(z) \leq E(c) \text{ for all } z \in Z$$

and equality holds if and only if $z = c$. This leads to a contradiction because $Z \setminus \{c\}$ is connected but, as Z is a representative in $H_{\text{ind}(\gamma)}(\mathcal{B}_x^{E(c)})$ that maps non-trivially to $H_{\text{ind}(\gamma)}(\mathcal{B}_x^{E(c)}, \mathcal{B}_x^{E(c)-})$, $Z \setminus \{c\}$ has to meet both components of $\mathcal{B}_x^{E(c)-}$ that are only connected by c . $\square$

Theorem 5.4. *The leaves of a singular Riemannian foliation $(M, \mathcal{F})$ with embedded leaves and without horizontal conjugate points are taut.*

Proof. At first, let N be regular. For each focal point x of N along a horizontal geodesic γ_x , we find a corresponding manifold K_x of dimension $\nu(x)$ within a neighboring leaf L that is specifically regular resp. of minimal type along γ_x as in the proof of theorem 5.1 and a global variation $f_x(s, t), s \in K_x$, of geodesics such that $f_x(s, 0) \in N, f_x(s, 1) = x$ for all $s \in K_x$ because of Proposition 4.13. Proposition 4.9 tells us that leaves of different dimension or different type than L are isolated along γ_x and therefore, we can carry over the whole proof of the previous theorem: Compactness of the image of K_x in N follows analogously as long as we choose

all these singular and exceptional leaves of higher type along γ_x to be centers of tubular neighborhoods of the covering. Hence, the only work to do regarding our Theorem 5.4 is an iteration of this idea. As in [BS58], we can use a fiber bundle construction.


The notation is the same as above. Denote the focal points on γ by

$$\{\gamma(t_j)\}_{j=1}^k, 1 = t_0 > t_1 > \dots > t_k > t_{k+1} = 0, \text{ with } \text{ind}(\gamma) = \sum_{j=1}^k \nu(\gamma(t_j)).$$

For all $j = 1, \dots, k$ let $f_j(s, \cdot) : [0, t_j] \rightarrow M$ denote the $\nu(\gamma(t_j))$ -dimensional variation of geodesics from N to $\gamma(t_j)$ parametrized by K_j (similar to the proof of Theorem 5.1). This variation consists of all minimal geodesics between K_j and $\gamma(t_j)$ continued up to the length $L(\gamma|_{[0, t_j]})$. In this situation holds

Lemma 5.5. *If $\gamma(t_i)$ is a focal point of N along γ , then $f_j(s, t_i)$ is a focal point of N of same multiplicity along $f_j(s, \cdot)$ for $j = 1, \dots, i - 1$ and for all $s \in K_j$. Moreover, we have*

$$N_{f_j(s, t)} = N_{\gamma(t)} \text{ for all } j = 1, \dots, k + 1 \text{ and } s \in K_j, t \in [0, t_j].$$



Proof. This results directly from Lemma 4.2 and Proposition 4.13 as well as from the variation formula of the energy. $\square$

Remark 5.6. As remarked in [Tö04], focal points along geodesics within such a variation are not fixed in general (i.e. in an arbitrary singular Riemannian foliation), neither are their multiplicities. Although easily proven with the help of Section 4 here, the lemma is essential to guarantee that the iterated bundle will be a manifold of dimension $\text{ind}(\gamma)$ as we will see now.

Knowing that along all geodesics of the variations f_j , there are the same focal points at the same time as along γ we can define

$$K := \{(s_1, \dots, s_k) \mid s_1 \in K_1(\gamma(1)), s_2 \in K_2(s_1), \dots, s_k \in K_k(s_{k-1})\},$$


where $K_i(s_{i-1})$ denotes the manifold of dimension $\nu(\gamma(t_i))$ which focalizes to the point t_i along the geodesic $f_{s_{i-1}}(t), t \in [0, t_{i-1}]$, for all i . And by the Lemma 5.5, K is a compact manifold, a fiber bundle, of dimension $\text{ind}(\gamma)$.

Now define the map $\lambda : K \rightarrow P(M, N \times p)$ as follows:

$$\lambda(s_1, \dots, s_k)(t) = \begin{cases} f_{s_i}(t), & t \in [t_i, t_{i+1}], i = 1, \dots, k \\ \gamma|_{[t_1, 1]}(t), & t \in [t_1, 1]. \end{cases}$$

The cycle $\lambda(K)$ thus consists of broken geodesics, all of same length; the only one that is non-broken and therefore the critical point of E_p is γ itself.

Sketch of the construction of $\lambda(K)$ in M .





Furthermore, $\lambda(K)$ has dimension $\text{ind}(\gamma)$ and is injective in the path space because each step of the iteration, i.e. of the bundle construction, yields only different broken geodesics. Hence, all requirements for a linking cycle for γ are fulfilled. By arbitrariness of γ , the claim follows for regular leaves N .

For a singular leaf N on γ , the argumentation is the same as above in the proof of Theorem 5.1: Assume p to be regular and construct the linking cycle of a regular leaf L near N . Then, add the geodesics connecting the endpoints of the cycle in L with N . The new manifold of broken geodesics fulfills all conditions for a linking cycle for γ . □

We will at last give an example in order to illustrate the cycle construction.

Example 5.7. Assume that the leaf N in $M = \mathbf{R}^3$ is a 2-torus, and the critical point in $P(M, N \times p)$ is a horizontal geodesic γ as in the figure on the right. The index of γ is two, because there are the two focal points p_1 and p_2 each of them with multiplicity one.



Let $K_2(s), s \in K$, be the appropriate, one-dimensional focal manifold, which can be identified as meridian through s of the torus.

Thus, the linking cycle of γ in $P(M, N \times p)$ is topologically the torus and consists of broken geodesics as outlined in this third figure.

The focal manifold K_1 can be identified with the outer latitude of the torus. There is an one-dimensional variation $f_s(t)$, $s \in K_1$, through geodesics starting perpendicularly in $s \in K_1 \subset N$ and ending in the focal point p_1 . On each of the geodesics lies exactly one focal point like p_2 on γ .




figure 3

REFERENCES

- [AT04] M. Alexandrino and D. Töben, Singular Riemannian Foliations on Simply Connected Spaces, *Differential Geom. Appl.*, 24 (2006), 383–397.
- [AT07] ———, Equifocality of a Singular Riemannian Foliation, *Proc. Amer. Math. Soc.*, 136 (2008) No. 9, 3271–3280.
- [Bo07] C. Boltner, On the Structure of Equidistant Foliations of Euclidian Space, arXiv: 0712.0245 [math.DG], 03 Dec 2007
- [BS58] R. Bott and H. Samelson, Applications of the Theory of Morse to Symmetric Spaces, *Amer. Jour. of Math.*, 80 (1958), 964–1029.
- [CC00] A. Candel and L. Conlon *Foliations*, volume 23 of *Graduate Studies in Mathematics*, AMS (2000).
- [CE75] J. Cheeger and D. G. Ebin, *Comparison Theorems in Riemannian Geometry*, North-Holland Publishing Company (1975).
- [dC93] M. P. do Carmo, *Riemannian Geometry, Mathematics: Theory and Applications*, Birkhäuser (1993).
- [GMPT] F. Giannoni, A. Masiello, P. Piccione and D. V. Tausk, A Generalized Index Theorem for Morse-Sturm Systems and Applications to Semi-Riemannian Geometry, *Asian J. Math.*, Vol. 5, No. 3 (2001), 441–472.
- [GTh02] C. Gorodski and G. Thorbergsson, Cycles of Bott-Samelson Type for Taut Representations, *Annals of Global Analysis and Geometry*, 21 (2002), 287–302.
- [GH91] K. Grove and S. Halperin, Elliptic Isometries, Condition (C) and Proper Maps, *Arch. Math.*, 56 (1991), 288–299.
- [HPT88] W.-Y. Hsiang, R. S. Palais and C.-L. Terng, The Topology of Isoparametric Submanifolds, *J. Differential Geometry*, 27 (1988), 423–460.
- [Ly07] A. Lytchak, Notes on the Jacobi Equation, *preprint* (2007) arXiv: 0708.2651 [math.DG].
- [Ly07a] ———, Geometric Resolution of Singular Riemannian Foliations, *preprint* (2007)
- [Ly07b] ———, Singular Riemannian Foliations on Spaces without Conjugate Points, *preprint* (2007)
- [LTh05] A. Lytchak and G. Thorbergsson, Variationally Complete Actions on Nonnegatively Curved Manifolds, *Illinois J. Math.*, 51 (2007), No. 2, 605–615.
- [LTh07] ———, Curvature Explosion in Quotients and Applications, *preprint* (2007).
- [Mi63] J. Milnor, *Morse Theory*, *Annals of Mathematics Studies*, No. 51, Princeton University Press 2nd Edition (1973).
- [Mol88] P. Molino, *Riemannian Foliations*, volume 73 of *Progress in Mathematics*, Birkhäuser (1988).
- [Mor34] M. Morse, *The Calculus of Variations in the Large*, volume 18 of *AMS Colloquium Publications*, AMS (1934).
- [PT89] R. S. Palais and C.-L. Terng, *Critical Point Theory and Submanifold Geometry*, volume 1353 of *Lecture Notes in Mathematics*, Springer Verlag (1989).
- [STh71] H. Seifert und W. Threlfall, *Variationsrechnung im Grossen*, Chelsea, New York (1971).
- [Ste74] P. Stefan, Accessible Sets, Orbits, and Foliations with Singularities, *Proc. London Math. Soc.*, 29 (1974), 699–713.
- [Su73] J. Sussmann, Orbits of Families of Vector Fields and Integrability of Distributions, *Transactions of the AMS*, 180 (1973), 171–188.

- [TTh97] ———, Taut Immersions into Complete Riemannian Manifolds, *Tight and Taut Submanifolds (Berkeley, CA, 1994)*, 181-228, *Math. Sci. Res. Inst. Publ.*, 32 Cambridge Univ. Press (1997).
- [Th83] G. Thorbergsson, Dupin Hypersurfaces, *Bull. London Math. Soc.*, 15 (1983), 493–498.
- [Tö04] D. Töben, Submanifolds with Parallel Focal Structure, *Doctoral thesis*, Universität zu Köln, 2004.
- [Wa81] R. Walter, Konvexität in Riemannschen Mannigfaltigkeiten, *Jahresbericht DMV*, Band 83, Teubner Verlag (1981), 1–31.
- [Wi09] S. Wiesendorf, *Doctoral thesis*, Universität zu Köln, in preparation.
- [Wi06] B. Wilking, A Duality Theorem for Riemannian Foliations in Nonnegative Sectional Curvature, *Geom. Funct. Anal.*, 17 (2007), No. 4, 1297–1320.

EVA NOWAK, MATHEMATISCHES INSTITUT, UNIVERSITÄT ZU KÖLN, WEYERTAL 86-90, 50931 KÖLN, GERMANY

E-mail address: enowak@mi.uni-koeln.de